

The sign of Δm_{31}^2 and the muon-charge asymmetry for atmospheric neutrinos

J. Bernabéu* and S. Palomares-Ruiz

*Departament de Física Teòrica
Universitat de València
46100 Burjassot, València, Spain*

Abstract

We discuss the possibility to measure the sign of Δm_{31}^2 from matter-induced charge asymmetries in atmospheric neutrino oscillations. The main conclusion is that an impact on the ν_μ survival probability requires the action of the MSW resonance, which becomes visible for baselines above ~ 7000 km.

*Talk given at the 7th International Workshop on Topics in Astroparticles and Underground Physics (TAUP2001), Laboratori Nazionali del Gran Sasso, Italy, September 8-12, 2001.

1 Introduction

Present evidence for neutrino masses and mixings can be summarized as: 1) the atmospheric $|\Delta m_{31}^2| \sim (1 - 5) \cdot 10^{-3} \text{eV}^2$ is associated with a mixing, θ_{23} , near to maximal [1]; 2) the solar Δm_{21}^2 prefers the LMA-MSW solution [2]; CHOOZ reactor data [3] give severe limits for $|U_{e3}|$. In this contribution we are going to discuss that contrary to a wide spread belief, Earth effects on the propagation of atmospheric neutrinos can become observable [4] even if $|U_{e3}|$ is small, but non-vanishing. This fact would allow to determine the sign of Δm_{31}^2 [5]. For baselines L smaller than the Earth diameter, appropriate for atmospheric neutrinos, $\frac{\Delta m_{21}^2}{4E} L \equiv \Delta_{21} \ll 1$, so that we will neglect the (1,2)-oscillating phase in vacuum against the (2,3)-one. This is a very good approximation, unless the high Δm_{21}^2 -region of the LMA solution turns out to be the solution to the solar problem. In that case we should take into account corrections of order $O(\frac{\Delta m_{21}^2}{\Delta m_{31}^2})$ (see eg. [6]).

In section 2 we discuss the correspondence between the determination of the sign of Δm_{31}^2 and the observation of the Earth effects in a transition involving ν_e . The change expected in the neutrino spectrum and mixing due to matter effects is pointed out. Section 3 studies the observability of the MSW-resonance, with a positive conclusion for baselines $L \gtrsim 7000$ km, and its impact on the survival probability, $\nu_\mu \rightarrow \nu_\mu$. Section 4 gives an analysis of the matter-induced CPT-odd asymmetry, together with the realistic charge-asymmetry expected for atmospheric neutrinos. In section 5 we present some conclusions.

2 The neutrino spectrum in matter

Current analyses leave us with two alternatives for the spectrum of the three active neutrino species, either hierarchical or degenerate.

The effective neutrino potential due to the charged current interaction of ν_e with the electrons in the medium is [7] $V \equiv \frac{a}{2E} = \sqrt{2}G_F N_e$, so that the effective hamiltonian, in the extreme relativistic limit, is given by [8]

$$H = \frac{1}{2E} \left\{ U \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \Delta m_{31}^2 \end{pmatrix} U^\dagger + \begin{pmatrix} a & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \right\} \quad (1)$$

In going from ν to $\bar{\nu}$, there are matter-induced CP- and CPT- odd effects associated with the change $a \rightarrow -a$. The additional change $U \rightarrow U^*$ is inoperative in the limit of (1). The effects we are going to discuss depend on the interference between the different flavors and on the relative sign between a and Δm_{31}^2 . As a consequence, an experimental distinction between the propagation of ν and $\bar{\nu}$ (the sign of a) will determine the sign of Δm_{31}^2 . An appreciable interference will be present if and only if there are appreciable matter effects. For atmospheric neutrinos, one needs the “connecting” mixing U_{e3} between the ν_e -flavor and the ν_3 mass eigenstate to show up.

For small s_{13} [3], even if the effects on the spectrum are expected to be small with respect to the decoupling of the ν_e -flavour in matter, there could be a substantial mixing

of ν_e with $\tilde{\nu}_3$ if one is near to a situation of level-crossing. This would lead to a resonant MSW behaviour [9].

$$\sin^2 2\tilde{\theta}_{13} = \frac{4 s_{13}^2 c_{13}^2}{(\alpha - \cos 2\theta_{13})^2 + 4 s_{13}^2 c_{13}^2} \quad , \quad \alpha \equiv \frac{a}{\Delta m_{31}^2} \quad (2)$$

But still $\langle \tilde{\nu}_1 | \nu_e \rangle = 0$, i. e., the ν_e has no overlap with the lowest mass eigenstate in matter. This vanishing mixing in matter is responsible for the absence of fundamental CP-violating effects, even if there are three non-degenerate mass eigenstates in matter. In vacuum, the absence of genuine CP-odd probabilities was due to the degeneracy $\Delta_{21} = 0$. The step from vanishing Δ_{21} in vacuum to the vanishing mixing U_{e1} in matter was termed a “transmutation” [10].

3 Observability of the MSW resonance

For atmospheric ν_μ neutrinos, matter effects in the survival probability $\nu_\mu \rightarrow \nu_\mu$ would be minute unless the resonance shows up. The resonance is not apparent even at $L = 3000$ km, appropriate for neutrino factories [11]. Is there a way out?

Again, a non-vanishing connecting mixing $s_{13} \neq 0$ provides the solution. Along with it, there is a resonance width which, when discussed in terms of the dimensionless parameter α , is given by

$$\alpha_R = \cos 2\theta_{13} \quad , \quad \Gamma_\alpha = 2 \sin 2\theta_{13} \quad (3)$$

One discovers that the oscillating phase on the resonance is non-vanishing, but given by the L -dependent relation

$$\tilde{\Delta}_{31(R)} = \Delta_{31} \frac{\Gamma_\alpha}{2} \quad (4)$$

If $L \ll L_{opt}$, with optimal L , L_{opt} , defined by $\tilde{\Delta}_{31(R)} = \pi/2$, the resonance does not affect the oscillation probability. On the contrary, around $L_{opt} = \frac{2\pi}{\tilde{a} \tan 2\theta_{13}}$, where $\tilde{a} = a/E$, the resonance becomes apparent and L_{opt} is independent of Δm_{31}^2 , which determines the resonant energy. For $L = L_{opt}$, the maximum mixing is accompanied by maximum oscillating factor.

Under these conditions, all channels would see the resonant effect. Contrary to non-resonant matter effects, the resonance only affects the (anti)neutrino channels if $\Delta m_{31}^2 > 0 (< 0)$.

4 Charge asymmetries

As discussed in section 3, matter effects distinguish neutrinos from antineutrinos. It is convenient to present them in terms of CP-odd (for appearance channels) and CPT-odd (for the survival probabilities) asymmetries. In the limit $\Delta_{21} = 0$, there is no room for




Figure 1: Upper panel: CPT-asymmetry, A_{CPT} , for different values of $\sin^2 2\theta_{13}$. From up to down: $\sin^2 2\theta_{13} = 0.005, 0.01, 0.05, 0.10, 0.16$ and $\Delta m_{31}^2 > 0$ as all the plots are symmetric with respect to the horizontal axis when $\Delta m_{31}^2 < 0$. Lower panel: Charge-asymmetry, A , for $\sin^2 2\theta_{13} = 0.05$ (dashed line) and 0.16 (solid line). The lower plots correspond to $\Delta m_{31}^2 > 0$ and the upper ones to $\Delta m_{31}^2 < 0$. For both panels, $L = 8000$ km, $\sin^2 2\theta_{23} = 1$ and $|\Delta m_{31}^2| = 3.2 \cdot 10^{-3} \text{ eV}^2$.

genuine CP violation. The interaction with matter will generate an asymmetry effect, however, which is not connected with the vacuum propagation.

For ν_μ and $\bar{\nu}_\mu$, one has

$$A_{CPT} = \frac{P(\nu_\mu \rightarrow \nu_\mu; L) - P(\bar{\nu}_\mu \rightarrow \bar{\nu}_\mu; L)}{P(\nu_\mu \rightarrow \nu_\mu; L) + P(\bar{\nu}_\mu \rightarrow \bar{\nu}_\mu; L)} \quad (5)$$

and it is represented in fig. 1 as function of the energy for a baseline of $L = 8000$ km and different values of $\sin^2 2\theta_{13}$. Around the resonance, A_{CPT} presents a plateau with non-vanishing appreciable values (depending on $\sin^2 2\theta_{13}$). The big asymmetries at 6 and 20 GeV correspond to low probabilities and they are not of interest. The negative (positive) asymmetry in the plateau is obtained for $\Delta m_{31}^2 > 0 (< 0)$. Obviously, it is symmetric with respect to the horizontal axis when changing the sign of Δm_{31}^2 . As we have seen above, the optimal baseline is inversely proportional to the θ_{13} mixing.

For atmospheric neutrinos, A_{CPT} cannot be separated out and the $\nu_e(\bar{\nu}_e)$ flux also contributes to the detection of $\nu_\mu(\bar{\nu}_\mu)$. Taking into account the CC cross-sections in the detector,

$$\begin{aligned} N(\mu^-; E) &= \sigma_{cc}(\nu_\mu) [\phi^o(\nu_\mu; E) P(\nu_\mu \rightarrow \nu_\mu) + \phi^o(\nu_e; E) P(\nu_e \rightarrow \nu_\mu)] \\ N(\mu^+; E) &= \sigma_{cc}(\bar{\nu}_\mu) [\phi^o(\bar{\nu}_\mu; E) P(\bar{\nu}_\mu \rightarrow \bar{\nu}_\mu) + \phi^o(\bar{\nu}_e; E) P(\bar{\nu}_e \rightarrow \bar{\nu}_\mu)] \end{aligned} \quad (6)$$

where $\phi^o(\nu_\mu; E)$ ($\phi^o(\bar{\nu}_\mu; E)$) and $\phi^o(\nu_e; E)$ ($\phi^o(\bar{\nu}_e; E)$) are the muon and electron (anti) neutrino fluxes, respectively, calculated from [12]. As in the important energy range, both cross-sections are, to good approximation, linear with the energy, one can build an asymmetry which eliminates what is induced by σ_{cc} in the form

$$A = \frac{N(\mu^-; E) - \frac{\sigma_{cc}(\nu_\mu)}{\sigma_{cc}(\bar{\nu}_\mu)} N(\mu^+; E)}{N(\mu^-; E) + \frac{\sigma_{cc}(\nu_\mu)}{\sigma_{cc}(\bar{\nu}_\mu)} N(\mu^+; E)} \quad (7)$$

In (7) there is still some asymmetry generated by the atmospheric neutrino fluxes. Contrary to A_{CPT} , the value of the muon-charge asymmetry is not symmetric with respect to the abscissa axis when changing the sign of Δm_{31}^2 . In fig. 1 we give the values of A for two values of $\sin^2 2\theta_{13}$. There is again an appreciable separation between the cases of positive and negative Δm_{31}^2 .

5 Conclusions

In the limit of $\frac{\Delta m_{21}^2}{4E}L \ll 1$, the main conclusions of this study are: i) The medium effects, which discriminate between neutrino and antineutrino propagation determine the sign of the atmospheric Δm_{31}^2 ; ii) for $s_{13} = 0$, electron neutrinos decouple from neutrino mixing in matter and have a definite effective mass in matter; iii) for $s_{13} \neq 0$, electron neutrinos mix with the third mass eigenstate neutrino and take part in the atmospheric neutrino oscillations; iv) electron neutrinos do not mix with the first mass eigenstate in matter, avoiding the generation of genuine CP-violating effects; v) non-resonant medium effects are already apparent in the sub-sominant channel $\nu_e \rightarrow \nu_\mu$ for baselines $L \sim 3000$ km, in both the mixing and the oscillation phase-shift; vi) the observation of matter effects in the ν_μ -survival probability requires the action of the MSW resonance, with baselines longer than $L \sim 7000$ km; vii) the optimal baseline depends on the value of s_{13} , but the effects are much cleaner in the region of the longest baselines without entering the Earth core [4] (nadir angles $\theta_n \gtrsim 33^\circ$). The corresponding muon-charge asymmetry shows an appreciable separation for the two possible signs of Δm_{31}^2 .

This work is supported by Grant AEN-99-0296.

References

- [1] H. Sobel, SuperKamiokande Collaboration, at Neutrino 2000, Sudbury, Canada (2000).
- [2] Y. Suzuki, SuperKamiokande Collaboration, at Neutrino 2000, Sudbury, Canada (2000); Q. R. Ahmad et al., SNO Collaboration, *Phys. Rev. Lett.* **87** (2001) 071301.
- [3] M. Apollonio et al., CHOOZ Collaboration, *Phys. Lett.* **B 420** (1998) 397.

- [4] M. C. Bañuls, G. Barenboim and J. Bernabéu, *Phys. Lett.* **B 513** (2001) 391, hep-ph/0102184; J. Bernabéu, A. Pérez, S. Palomares-Ruiz and S. T. Petcov, hep-ph/0110071.
- [5] V. Barger, S. Geer, R. Raja and K. Whisnant, *Phys. Rev.* **D 62** (2000) 013004, hep-ph/9911524.
- [6] I. Mocioiu and R. Schrock, *J. High Energy Phys.* **0111** (2001) 050, hep-ph/0106139; M. Freund, *Phys. Rev.* **D 64** (2001) 053003, hep-ph/0103300.
- [7] L. Wolfenstein, *Phys. Rev.* **D 17** (1978) 2369.
- [8] T. K. Kuo and J. Pantaleone, *Rev. Mod. Phys.* **61** (1989) 937.
- [9] S. P. Mikheyev and A. Yu. Smirnov, *Sov. J. Nucl. Phys.* **42** (1985) 913.
- [10] J. Bernabéu and M. C. Bañuls, *Nucl. Phys.* **87 (Proc. Suppl.)** (2000) 315, hep-ph/0003299.
- [11] A. De Rújula, M. B. Gavela and P. Hernández, *Nucl. Phys.* **B 547** (1999) 21, hep-ph/9911390; K. Dick, M. Freund, M. Lindner and A. Romanino, *Nucl. Phys.* **B 562** (1999) 29, hep-ph/9903308.
- [12] G. Fiorentini, V.A. Naumov and F.L. Villante, *Phys. Lett.* **B 510** (2001) 173, hep-ph/0103322.